



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

23rd International Grassland Congress

Genetic Diversity and Relationship among 19 Accessions of Sorghum-Sudangrass Hybrids as Revealed by SSR Markers

Qiuwen Zhan

Anhui Science and Technology University, China

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/23/4-1-1/5>

The 23rd International Grassland Congress (Sustainable use of Grassland Resources for Forage Production, Biodiversity and Environmental Protection) took place in New Delhi, India from November 20 through November 24, 2015.

Proceedings Editors: M. M. Roy, D. R. Malaviya, V. K. Yadav, Tejveer Singh, R. P. Sah, D. Vijay, and A. Radhakrishna

Published by Range Management Society of India

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Genetic diversity and relationship among 19 accessions of Sorghum-sudangrass hybrids as revealed by SSR markers

Qiuwen Zhan

Anhui Science and Technology University, Fengyang, China

Corresponding author e-mail: qwzhan@163.com

Keywords: Cluster analysis, Genetic diversity, Relationship, Sorghum-sudangrass hybrids, SSR

Introduction

Sorghum (*Sorghum bicolor*) and sudangrass (*S. sudanense*) are two members of the agriculturally important genus *Sorghum*. Hybrids of these two species have exhibited favorable forage yields and overall improved quality and disease resistance (Zhan and Qian, 2004). Sorghum-sudangrass hybrids (*Sorghum bicolor* × *S. sudanense*) are used as a forage crop and recommended as an alternative to corn in USA (Kilcer *et al.*, 2003). In recent years, the planting areas of sorghum-sudangrass hybrids have increased in China. SSR has been widely employed in crop germplasm identification, genetic mapping and diversity analysis (Zhan *et al.*, 2008, Li *et al.*, 2010). However, no reports have been published on studies of the genetic diversity of main sorghum-sudangrass hybrids in China. The objective of this current study was to evaluate the genetic diversity and relationship among 19 accessions of sorghum-sudangrass hybrids as revealed by SSR markers.

Materials and Methods

19 accessions of sorghum-sudangrass hybrids and one accession from each of three related species (*S. sorghum*, *S. Sudanense*, and *Zea mays*) were included in the study (Table 1). DNA extraction and SSR PCR reactions were performed as described (Zhan *et al.*, 2008). A molecular marker data matrix was established by recording 1 for the presence of a specific band and 0 for the absence of the band. Statistical analysis software, NTSYS-pc, was used, in which the measure of similarity coefficient was Dice (Rohlf, 2000). The UPGMA method was used for cluster analysis and phylogenetic tree construction. Gene diversity was evaluated with the polymorphic information content (PIC) (Zhan *et al.*, 2008).

Table 1: The name, type and source of 22 accessions tested

NO.	Name	Type	Source
1	Everlush	<i>S. bicolor</i> × <i>S. sudanense</i>	Australia
2	MMR327/36	<i>S. bicolor</i> × <i>S. sudanense</i>	USA
3	MMR327/38	<i>S. bicolor</i> × <i>S. sudanense</i>	USA
4	Verdeo-45	<i>S. bicolor</i> × <i>S. sudanense</i>	Argentina
5	MMR352/40	<i>S. bicolor</i> × <i>S. sudanense</i>	USA
6	Pacesetter	<i>S. bicolor</i> × <i>S. sudanense</i>	USA
7	Sweet N Honey II	<i>S. bicolor</i> × <i>S. sudanense</i>	USA
8	Evergreen	<i>S. bicolor</i> × <i>S. sudanense</i>	USA
9	Pioneer	<i>S. bicolor</i> × <i>S. sudanense</i>	USA
10	Wancao No.3	<i>S. bicolor</i> × <i>S. sudanense</i>	Anhui,China
11	Jincao No.5	<i>S. bicolor</i> × <i>S. sudanense</i>	Shanxi,China
12	Jincao No.1	<i>S. bicolor</i> × <i>S. sudanense</i>	Shanxi,China
13	Jicao No.2	<i>S. bicolor</i> × <i>S. sudanense</i>	Hebei,China
14	S.Africa-Su	<i>S. bicolor</i> × <i>S. sudanense</i>	South Africa
15	Wancao No.2	<i>S. bicolor</i> × <i>S. sudanense</i>	Anhui,China
16	Jincao No.6	<i>S. bicolor</i> × <i>S. sudanense</i>	Shanxi,China
17	Jicao No.1	<i>S. bicolor</i> × <i>S. sudanense</i>	Hebei,China
18	Jicao No.3	<i>S. bicolor</i> × <i>S. sudanense</i>	Hebei,China
19	BMR1	<i>S. bicolor</i> × <i>S. sudanense</i>	Shanxi,China
20	TX623A	<i>S. sorghum</i>	USA
21	S722	<i>S. sudanense</i>	India
22	Yunyou 199	<i>Z. mays</i>	Yunnan,China

Results and Discussion

The 79 SSR primer pairs generated a total of 259 polymorphic alleles, with an average of 3.28 alleles per locus, and a range of 1 to 7 alleles. The polymorphic rate was 91.89% within 22 accessions, and the average genetic diversity (PIC) value was 0.440. Except three relatives (sorghum, sudangrass and maize), a total of 222 polymorphic alleles were generated within 19 sorghum-sudangrass hybrids, with a polymorphism rate of 79.73%. The SSR profile generated by primer *astu224* was shown in Figure 1. The PIC value was 0.421. Genetic similarity (GS) values between all varieties ranged from 0.376 to 0.923 and were used to produce a dendrogram (Figure 2). The 22 accessions were clustered into three groups (GS=0.696), specifically, group I (sorghum-sudangrass hybrids), group II (sorghum and sudangrass), group III (maize). Results of our analyses suggest the sudangrass/sorghum relationship is sufficiently close, but the hybrids reveal obvious heterosis. Therefore, all accessions of sorghum-sudangrass hybrids were clustered into one group. GS values between 19 sorghum-sudangrass hybrids ranged from 0.646 to 0.923.




Fig 1: The amplification results of primer *astu224* in the 22 cultivars




Fig 2: The clustering result of 22 cultivars based on SSR coefficient

Conclusion

Except three relatives (sorghum, sudangrass and maize), a total of 222 polymorphic alleles were generated within 19 sorghum-sudangrass hybrids, with a polymorphism rate of 79.73%. The PIC value was 0.421. These varieties from the same region, especially those from the same breeding institution, exhibited small genetic variations and high genetic similarities. At present, more attention should be paid to discovery and innovation in the breeding of sorghum-sudangrass hybrids.

References

- Kilcer, T. F., Q. M. Ketterings, P. Cerosaletti, P. Barney and J. H. Cherney. 2003. Cutting Management for Brown Mid Rib Sorghum Sudangrass. What's Cropping Up? *Newsletter*, 13(4): 4-6.
- Li, J. Q., L. H. Wang, Q. W. Zhan and J. C. Fan. 2010. Genetic analysis and mapping with SSR markers of the brown midrib gene bmr-6 in *Sorghum bicolor*. *Acta Prataculturae Sinica*, 19(5): 273-277.
- Rohlf, F. J. 2000. *NTSYS-pc, Numerical Taxonomy and Multivariate Analysis System*, Version 2.10e. Applied Biostatistics Inc., New York.
- Zhan, Q. W., T. Z. Zhang, B. H. Wang and J. Q. Li. 2008. Diversity Comparison and Phylogenetic Relationships of *S. bicolor* and *S. sudanense* as Revealed by SSR Markers. *Plant Science* 174 (1): 9-16.
- Zhan, Q.W. and Z. Q. Qian. 2004. Heterosis Utilization of Hybrid between Sorghum [*Sorghum bicolor* (L.) Moench] and Sudangrass [*Sorghum sudanense* (Piper) Stapf]. *Acta Agronomica Sinica*, **30(1)**:73-77.

Acknowledgement

This research was supported by the National Natural Science Foundation of China (31301383, 31071470), the Key-construction Subject Plan of Anhui Province (WanJiaoMiKe [2014]28) and the subject of the twelfth five-year-plan in national science and technology for the rural development in China (2011BAD17B03).